arXiv:hep-ph/9809434v3 11 Jan 1999

# THE EIKONAL APPROACH TO CALCULATION OF THE MULTIPHOTON EXCHANGE CONTRIBUTIONS TO THE TOTAL CROSS SECTIONS OF $\pi^+\pi^-$ ATOM INTERACTION WITH ORDINARY ATOMS

L.AFANASYEV and O.VOSKRESENSKAYA
*Joint Institute for Nuclear Research,*
*141980 Dubna, Moscow Region,*
*Russia*

## Abstract

The total cross sections of the relativistic $\pi^+\pi^-$ atom interactions with ordinary atoms are obtained in the eikonal approach which takes into consideration all multiphoton exchange processes. Contribution of these processes strongly depends on the atom nucleus charge $Z$ and varies from 1.5% for Titanium ($Z = 22$) to 14% for Tantalum ($Z = 73$)

An accurate measurement of the $\pi^+\pi^-$ atom (dimesoatom) lifetime in the experiment DIRAC [1] will allow one to check a high precision prediction of the Chiral Perturbation Theory. An interaction of the $\pi^+\pi^-$ atoms with ordinary atoms is an essential part of this experiment, as the atom observation bases on their breakup (ionization) while passing through the target where they are produced.

Cross sections of $\pi^+\pi^-$ atoms with ordinary atoms were usually calculated in the first Born approximation. In the paper [2] the cross sections for few low states of dimesoatom were calculated in Coulomb-modified Glauber approximation. It has been shown that the multiphoton exchange processes play a significant role in the interaction of $\pi^+\pi^-$ atoms with atoms.

In the Glauber approximation the total cross sections of the coherent interaction of $\pi^+\pi^-$ atom with ordinary atoms could be written [2] as:

$$\sigma^{tot}_{nlm} = 2\,\mathrm{Re}\int d^2b\, d^3r |\psi_{nlm}(\vec{r})|^2 \left[1 - \exp\left(i\chi(\vec{b} - \vec{s}/2) - i\chi(\vec{b} + \vec{s}/2)\right)\right] . \qquad (1)$$

Here $\vec{s} = \vec{r}_\perp$ is the projection of the vector $\vec{r}$ on the plane of the impact parameter $\vec{b}$, $\psi_{nlm}(\vec{r})$ is the wave function of $\pi^+\pi^-$ atom in the state with principal, orbital and magnetic quantum numbers $n$, $l$ and $m$, respectively. The phase shift $\chi(\vec{b})$ is expressed via the screened Coulomb potential of the target atom:

$$\chi(\vec{b}) = \frac{1}{v}\int\limits_{-\infty}^{\infty} U(\sqrt{b^2 + z^2}\,)\, dz\, . \qquad (2)$$

Here $v$ is the velocity of dimesoatom in the lab frame.

Let us write (1) in another form taking into account the following relations:

$$\begin{aligned} \exp\left(i\chi(\vec{b})\right) &= 1-\gamma(b)\,, & (3)\\ \gamma(b) &= \frac{1}{2\pi i}\int f(q)\exp\left(-iqb\right)d^2q\,, & (4)\\ f(q) &= \frac{i}{2\pi}\int\left[1-\exp\left(i\chi(\vec{b})\right)\right]\,\exp\left(i\vec{q}\vec{b}\right)d^2b & \\ &= i\int\limits_0^\infty\left[1-\exp\left(i\chi(\vec{b})\right)\right]\,J_0(qb)b\,db\,. & (5) \end{aligned}$$

Here $f(q)$ is the amplitude of the elastic Coulomb $\pi A$-scattering normalized by the relations:

$$\begin{aligned} \sigma_{\pi A}^{tot} &= 4\pi\,\mathrm{Im}\,f(0)\,, & (6)\\ \frac{d\,\sigma_{\pi A}}{d\vec{q}} &= |f(q)|^2\,. & (7) \end{aligned}$$

Then it is easy to get the total cross section in the form

$$\begin{aligned} \sigma_{nlm}^{tot} &= 2\int|f(q)|^2\left[1-S_{nlm}(\vec{q})\right]\,d^2q\,, & (8)\\ S_{nlm}(\vec{q}) &= \int|\psi_{nlm}(\vec{r})|^2\,\exp\left(i\vec{q}\vec{r}\right)d^3r\,. & (9) \end{aligned}$$

Here $S_{nlm}(\vec{q})$ is the elastic form factor of the $\pi^+\pi^-$ atom in the state with the quantum numbers $n$, $l$ and $m$.

The expressions (8) and (9) together with the results obtained in the paper [3] for the transition form factors of the hydrogen-like atoms allow one to calculate the total cross sections for any state of the dimesoatom. However, in this paper we only consider the cross sections averaged over the magnet quantum number:

$$\sigma_{nl}^{tot}=\frac{1}{2l+1}\sum_m\sigma_{nlm}^{tot}\,. \tag{10}$$

The wave function of $\pi^+\pi^-$ can be written as a product of the radial and angular parts:

$$\psi_{nlm}(\vec{r})=R_{nl}(r)\,Y_{lm}(\Theta,\phi)\,. \tag{11}$$

Taking into account the normalization

$$\frac{1}{2l+1}\sum_m|Y_{lm}(\Theta,\phi)|^2=1\,, \tag{12}$$

we have

$$\sigma_{nl}^{tot}=4\pi\int|f(q)|^2\left(1-S_{nl}(q)\right)\,q\,dq\,, \tag{13}$$

$$\begin{aligned} S_{nl}(q) &= \frac{1}{2l+1}\sum_m S_{nlm}(q)=\int|R_{nl}|^2\exp\left(i\vec{q}\vec{r}\right)d^3r=4\pi\int\limits_0^\infty|R_{nl}|^2\,\frac{\sin qr}{q}\,r\,dr & \\ &= \frac{4\pi}{q}\,\mathrm{Im}\int\limits_0^\infty|R_{nl}|^2\exp\left(iqr\right)r\,dr\,. & (14) \end{aligned}$$

The radial wave function of the hydrogen-like atom is expressed in terms of the Laguerre polynomial. So integration in (13) is reduced to the hypergeometric functions using the expression [4]

$$\int_0^\infty e^{-bx}\, x^\alpha\, L_n^\alpha(\lambda x)\, L_m^\alpha(\mu x) =$$

$$\frac{\Gamma(m+n+\alpha+1)}{\Gamma(m+1)\Gamma(n+1)}\, \frac{(b-\lambda)^n (b-\mu)^m}{b^{n+m+\alpha+1}} F\left[-m, -n, -m-n-\alpha; \frac{b(b-\lambda-\mu)}{(b-\mu)(b-\lambda)}\right] \tag{15}$$

Finally for the form factor we have

$$S_{nl}(q) = \frac{\sin 2n\phi\, (\cos\phi)^{2l+4}}{n\, \sin 2\phi}\, F(l+1-n, l+1+n; 1; \sin^2\phi)\,, \tag{16}$$

$$\phi = \arctan\left(\frac{nq}{\alpha m_\pi}\right).$$

For the numerical calculation we use the Moliére parametrization of the Thomas-Fermi potential [5]

$$U(r) = Z\alpha \sum_{i=1}^{3} \frac{c_i e^{-\lambda_i r}}{r}\,; \tag{17}$$

which allows one to obtain the exact expression for the phase shift $\chi(b)$

$$\chi(b) = \frac{2Z\alpha}{v} \sum_{i=1}^{3} c_i\, K_0(b\lambda_i)\,, \tag{18}$$

$$c_1 = 0.35, \quad c_2 = 0.55, \quad c_3 = 0.1\,;$$

$$\lambda_1 = 0.3\lambda_0, \quad \lambda_2 = 1.2\lambda_0, \quad \lambda_3 = 6\lambda_0, \qquad \lambda_0 = m_e \alpha Z^{1/3}/0.885\,.$$

The numerical results were calculated for the Titanium ($Z = 22$) and Tantalum ($Z = 73$) targets. The velocity of $\pi^+\pi^-$ atom was taken as $v = 1$. Figures 1–4 show the most important dependencies. We can conclude that the contribution of the multiphoton exchange processes strongly depends on the target atom nucleus charge $Z$ and varies from 1.5% for Titanium to 14% for Tantalum.

In the DIRAC experiment the dimesoatom lifetime is going to be measured for various targets with an accuracy of 10% using the breakup probability [1] which is calculated base on the interaction cross sections. So account of the multiphoton processes is essential for an interpretation of the experiment results.

Authors would like to thank A.V.Tarasov from JINR, professor J.Hüfner from Heidelberg Universität and professor B.Kopeliovich from Max-Plank-Institut für Kernphysik for helpful discussions. This work is partially supported by RFBR grant 97–02–17612.



Figure 1: Total cross sections of $\pi^+\pi^-$-atom interactions in $n$S states with Tantalum ($Z = 73$) versus the principal quantum number $n$ in the Born and Glauber approximation.



Figure 2: Relative difference between the total cross sections of $\pi^+\pi^-$-atom interaction in $n$S states with Tantalum ($Z = 73$) calculated in the Born and Glauber approximation versus the principal quantum number $n$.



Figure 3: Relative difference between the total cross sections of $\pi^+\pi^-$-atom interaction in $n$S states with Titanium ($Z = 22$) calculated in the Born and Glauber approximation versus the principal quantum number $n$.



Figure 4: Dependence of the total cross sections of $\pi^+\pi^-$-atom interaction with Tantalum on the orbital quantum number $l$ for the various principal quantum number $n$.

# References

[1] Adeva B. et al., *Lifetime measurement of* $\pi^+\pi^-$ *atoms to test low energy QCD predictions*, Proposal to the SPSLC, CERN/SPSLC 95–1, SPSLC/P 284, Geneva, 1995.

[2] Tarasov A.V. and Christova I.U., *JINR Communication* P2-91-10, Dubna, 1991.

[3] Afanasyev L.G. and Tarasov A.V., *Yad. Fiz.* **59** (1996) 2212.

[4] Gradshtein I.S. and Ryzhik I.M., *Tables of Intergals, Series and Products*, (5th edition, Nauka Publication, Moscow, 1971).

[5] Molière G., *Z. Naturforsch.* **2A** (1947) 3.